

For Clear Contrast

	Diatrizoic Acid Meglumine Salt % w/v	Diatrizoic Acid Sodium Salt % w/v	рН	Viscosity at 37°C	lodine concentration mg / ml
Trazograf 76% (Inj.)	66	10	6.0 to 7.7	8.4	370
Trazograf 60% (Inj.)	60	-	6.0 to 7.7	4.1	282
Trazogastro Oral	66	10	6.0 to 7.7	-	370

INDICATIONS:

Trazograf 76%

- I.V. Urography
- Computerised Tomography
- Aortography
- Angiography
- Urethrography
- Digital Subtraction Angiography
- Angiocardiography

Trazograf 60%

- Angiography (Cerebral, Visceral, Peripheral)
- Phlebography
- Digital Subtraction Angiography
- Computerised Tomography
- Hysterosalpingography
- Arthrography

Trazogastro

- Computerised Tomography for abdomen
- Visualisation of the Gastro-Intestinal Tract

CONTRAINDICATIONS & RISKS:

Manifest hyperthyroidism, decompensated cardiac Insufficency. Hysterosalpingography must not be performed during pregnancy & in the presence of acute inflammatory process in the pelvic cavity. In Hypersensitivity to iodinated contrast media, severe

Impairment of hepatic or renal function, cardiac & circulatory insufficiency, pulmonaryemphysema, poor general health, Cerebral arteriosclerosis, juveline type diabetes or diabetes of long standing duration, cerebral spasmodic condition, latent hyperthyroidism, gland nodular goitre, multiple Myeloma, polyuria, oliguria, gout in infants and young children the need of examination particularly needs careful considerations, pre-medication with alpha-receptor blockers is recommended in phaechromocytoma patients because of the risk of hypertensive crisis.

PLEASE NOTE:

Details on tolerance and side effects, individual hypersensitivity, adminstration during pregnancy and in patients with allergic disposition, subsequent diagnosis of a thyroid disease with iodine isotopes, special notes as well as suggestions for the treatment of contrast medium incidents are contained in the package insert.

PRESENTATION:

TRAZOGRAF 76%:

100 ml Bottle, 50ml Vial, 20 ml Ampoule

TRAZOGRAF 60 %:

50ml Vial, 20ml Ampoule

TRAZOGASTRO:

100ml Bottle, 30ml Bottle



* Registered Trademark

lanufactured under Technical Collaboration

rith JUSTESA IMAGEN SA 28028 Madrid, Spain

Further information available on request from

B. CHEMICALS & PHARMACEUTICALS LIMITED

Cnergy IT Park, Unit A2, 3rd Floor, Unit A, 8th Floor, Appa Saheb Marathe Marg, Prabhadevi Mumbai 400 025, Phone: +91 22 2439 5200 / 2439 5500 Website: www.ibcgl.com

Unique's

TRAZOGRAF*

For Clear Contrast



TRAZOGRAF* 76%



TRAZOGRAF* 60%



- The physiological Na Conc.(360 mg / 100 mL)
- The Right Ratio of Meglumine : Sodium (66 : 10)
- Stabilizer CaNa, EDTA

- Least effect on the electrolyte balance
- Optimal cardiac tolerance
 - Reduced risk of Ventricular fibrillation

No Sodium

Low Viscosity



COMPOSITION

Diatrizoic Acid Salts

Meglumine Salt 66% w/v
Sodium Salt 10% w/v

IODINE

CONCENTRATION: 370 mg/ ml

 20 mL Ampoule
 7.4 g

 50mL Vial
 18.5 g

 100 mL Inf. Bottle
 37.0 g





pH: 6.0 to 7.7 VISCOSITY (mPa.s or cP)

at 37°C 8.4

INDICATIONS

- I.V. Urography
- Computerised Tomography
- Aortography
- Angiography
- Urethrography
- Digital Subtraction Angiography
- Angiocardiography

COMPOSITION

Diatrizoic Acid Salts

Meglumine Salt 60% w/v

IODINE

CONCENTRATION: 282 mg/ ml

20 mL Ampoule 5.64 g 50mL Vial 14.1 g



pH: 6.0 to 7.7

VISCOSITY (mPa.s or cP)

at 37°C 4.1

INDICATIONS

- Angiography (Cerebral, Visceral, Peripheral)
- Phlebography
- Digital Subtraction Angiography
- Computerised Tomography
- Hysterosalpingography
- Arthography

TRAZOGASTRO*

Only for Oral Use **COMPOSITION**

Diatrizoic Acid Salts

in Flavoured aqueous base

IODINE

CONCENTRATION: 370 mg/ ml

pH: 6.0 to 7.7



Easy to use - convenient



Economical

INDICATIONS

- Computerised Tomography for abdomen
- Visualisation of the Gastro-Intestinal Tract

Standard formula with wetting and flavouring agent

Better patient acceptance - Orange Flavour