

Sodium Alginate and Potassium Bicarbonate Chewable Tablets (500mg+100mg)

RANRAFT®
TABLET

GENERIC NAME

Sodium Alginate & Potassium Bicarbonate Chewable Tablets (500mg + 100mg)

QUALITATIVE AND QUANTITATIVE COMPOSITION

Each uncoated chewable tablet contains:
Sodium Alginate IP 500 mg
Potassium Bicarbonate BP 100 mg
Colour: Quinoline Yellow

DOSAGE FORM AND STRENGTH

Uncoated Chewable Tablet

CLINICAL PARTICULARS**THERAPEUTIC INDICATION**

Treatment of symptoms resulting from the reflux of acid, bile and pepsin into the oesophagus such as acid regurgitation, heartburn, indigestion (occurring due to the reflux of stomach contents), for instance, after gastric surgery, as a result of hiatus hernia, during pregnancy, accompanying reflux oesophagitis, including symptoms of laryngopharyngeal reflux such as hoarseness and other voice disorders, sore throats and cough. It can also be used to treat the symptoms of gastro-oesophageal reflux during concomitant treatment with or following withdrawal of acid suppressing therapy.

Posology and method of administration

Adults and children 12 years and over: One to two tablets after meals and at bedtime.

Children under 12 years: Should be given only on medical advice.

Elderly: No dose modifications necessary for this age group.

Hepatic Impairment: No dose modification necessary.

Renal Insufficiency: Caution if highly restricted salt diet is necessary

Method of administration: For oral administration only

The tablet must be chewed thoroughly before swallowing

CONTRAINDICATIONS

This medicinal product is contraindicated in patients with known or suspected hypersensitivity to the active substances or to any of the excipients.

SPECIAL WARNINGS AND PRECAUTIONS FOR USE

If symptoms do not improve after 7 days, the clinical situation should be reviewed

This medicinal product contains 53.22 mg sodium per tablet, equivalent to 2.7% of the WHO recommended maximum daily intake for sodium.

DRUG INTERACTION

A time-interval of 2 hours should be considered between this formulation intake and the administration of other medicinal products, especially tetracyclines, fluoroquinolones, iron salts, thyroid hormones, chloroquine, bisphosphonates, and estramustine. Alcohol consumption makes the stomach produce more acid, which can further lead to heartburn. Therefore, avoid alcohol consumption.

USE IN SPECIAL POPULATION**Fertility, pregnancy and lactation Pregnancy:**

Clinical studies in more than 500 pregnant women as well as a large amount of data from post-marketing experience indicate no malformative nor foeto/neonatal toxicity of the active substances. This drug can be used during pregnancy, if clinically needed.

Breast feeding:

No known effect on breast fed infants. This drug can be used during breast feeding.

Fertility:

No known effect on human fertility

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

None

UNDESIRABLE EFFECTS

Adverse reactions have been ranked under headings of frequency using the following convention: very common (1/10), common (1/100 and < 1/10,000) uncommon (1/1000 and 1/100), rare (1/10,000 and < 1/1000), very rare (<1/10,000) and not known (cannot be estimated from the available data).

System Organ Class	Frequency	Adverse Event
Immune System Disorders	Very rare	Anaphylactic and anaphylactoid reactions. Hypersensitivity reactions such as urticaria.
Respiratory, Thoracic and Mediastinal Disorders	Very rare	Respiratory effects such as bronchospasm
Gastrointestinal Disorders	Uncommon	Diarrhoea, nausea, vomiting.

Overdose**Symptoms:**

Symptoms are likely to be minor; some abdominal discomfort may be experienced.

Management:

In the event of overdose symptomatic treatment should be given

PHARMACOLOGICAL PROPERTIES**Mechanism of Action**

The mode of action of proposed FDC is local and does not depend on absorption into the systemic circulation.

Pharmacodynamic properties

Pharmacotherapeutic classification: A02BX 13. Other drugs for peptic ulcer and gastro-oesophageal reflux disease.

On ingestion the tablet reacts with gastric acid to rapidly form a raft of alginic acid gel having near-neutral pH which floats on the stomach contents effectively impeding gastro-oesophageal reflux for up to 4 hours, and protecting the oesophagus from acid, pepsin and bile. In severe cases the raft itself may be refluxed into the oesophagus in preference to the stomach contents and exert a demulcent effect. In addition in vitro evidence has shown that the raft has a secondary action and is able to entrap bile and pepsin within its structure, further protecting the oesophagus from these gastric components.

Pharmacokinetic properties

The mode of action of Sodium alginate & Potassium bicarbonate tablets is physical and does not depend on absorption into the systemic circulation.

NON CLINICAL PROPERTIES**Animal toxicology or Pharmacology**

No preclinical findings relevant to the prescriber have been reported

DESCRIPTION


This product (chewable tablet) contains sodium alginate & bicarbonate and it is Indicated for the treatment of heartburn and indigestion.

Sodium Alginate:

Chemical Name: sodium 3,4,5,6-tetrahydroxyoxane-2-carboxylate Molecular

Formula: (C₆H₇NaO₆)_n

Structural Formula:


**Potassium bicarbonate:**

Chemical Name: Potassium hydrogencarbonate

Molecular Formula: KHCO₃

Molecular weight: 100.115 g/mol

Structural Formula:

**PHARMACEUTICAL PARTICULARS****Incompatibility****Shelf Life**

Refer carton

Packaging Information

Blister of 10 tablets

Storage instructions

Store protected from moisture, at a temperature below 30oC.

PATIENT COUNSELLING INFORMATION

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.



Marketed by & © Regd. Trade Mark of:
J. B. CHEMICALS & PHARMACEUTICALS LTD.
Neelam Centre, 'B' Wing, Hind Cycle Road,
Worli, Mumbai - 400 030, India.

Note: This prescribing information is applicable for India Market only.

DATE OF REVISION: Feb 2025